

Genital Scabies

Skin Care & Health
skin-care-health.org/genital-scabies.html

Scabies

Scabies is caused by tiny parasitic mites. They are smaller than a pinhead and burrow into the skin and lay eggs. A more severe and uncommon form of the condition occurs when there are many mites in the skin. This is called crusted scabies, and

How do you get scabies?	3
What are the signs and symptoms?	3
If I think I have scabies, how soon can I have a check-up?.....	4

Scabies

How do you get scabies?

Scabies can be passed from one person to another by close body or sexual contact with someone who has scabies.

The mites which cause scabies can be found in the genital area, on the hands, between the fingers, on the wrists and elbows, underneath the arms, on the abdomen, on the breasts, around the nipples in women, on the feet and ankles, and around the buttocks.

The mites can live for up to 72 hours off the body, so it is possible for scabies to be spread by clothing, bedding and towels.

What are the signs and symptoms?

Some people will not have any visible signs or symptoms at all, or may not be aware of them. It can take up to six weeks after coming into contact with scabies before signs and symptoms appear. Signs and symptoms are the same for both men and women. You might notice:

- Intense itching in the affected areas which may only be noticed at night, or which becomes worse in bed at night or after a hot bath or shower.
- An itchy red rash or tiny spots. Sometimes the diagnosis can be difficult because the rash can look like other itchy conditions, such as eczema.
- Inflammation or raw, broken skin in the affected areas – usually caused by scratching.

Scabies mites are very tiny and impossible to see with the naked eye. Fine silvery lines are sometimes visible in the skin where mites have burrowed.

Sometimes scabies will be noticed during a

routine genital or medical examination even if a doctor or nurse isn't looking for it.

If I think I have scabies, how soon can I have a check-up?

You can only be certain you have scabies if you have a check-up. You can have a check-up as soon as you have signs or symptoms for the doctor or nurse to look at, or if you think you might have been in contact with scabies. Some people won't develop visible signs or symptoms straightaway, and you may be asked to come back for another appointment later on.

What does the check-up involve?

A doctor or nurse can often tell if you have scabies just by looking at the affected areas. They may gently take a skin flake from one of the areas and look at it under a microscope to see if there is a mite present.

In some cases, treatment will be suggested if scabies is suspected, even if it cannot be confirmed.

Where can I get a check-up?

There are a number of services you can go to. Choose the service you feel most comfortable with.

A test can be done at:

- a genitourinary medicine (GUM) or sexual health clinic
- your general practice; ask your doctor or practice nurse
- some contraception clinics and young people's services.

If you know you have scabies you can go to a pharmacy for advice and treatment.

What is the treatment for scabies?

- Treatment is simple and involves using a special cream or lotion. The doctor, nurse or pharmacist will advise you on what treatment to use and how to use it.
- You apply the cream or lotion usually to the whole body from the neck downwards. This ideally should be done overnight. The treatment should be rinsed off after 12 hours.
- You should wash clothing, bedding and towels in a washing machine on a very hot cycle (50°C or higher) to kill the mites and avoid re-infection.
- You can also buy treatments for scabies from pharmacies. These are useful for anyone who is sure they have scabies and wants to self-treat. The pharmacist will be able to advise if you have any questions, or are unsure how to use the treatment.
- If you decide to treat yourself, you may still want to consider having a sexual health check, to make sure you don't have a sexually transmitted infection.
- Do tell the doctor, nurse or pharmacist if you are, or think you might be, pregnant or if you are breastfeeding. This will affect the type of treatment you are given.
- Close contacts in your household should be treated at the same time, as well as your sexual partner, even if they do not have any signs or symptoms.
- There is no evidence that complementary therapies can cure scabies.

When will the signs and symptoms go away?

If you use the treatment according to the instructions it is rare for it not to work.

Even after successful treatment, the itching or rash may continue for a few weeks. Special tablets or creams (antihistamines), or anti-irritant lotions such as calamine, can ease the itching.

How soon can I have sex again?

It is strongly advised that you do not have any sexual intercourse, including vaginal, anal or oral sex, or close body contact, until you **and** your partner have both finished the treatment and any follow-up treatment. This is to help prevent you being reinfected or passing the infection on to someone else.

Do I need to have a check-up to see if the scabies has gone?

You may need to go back to check the scabies has gone away and that you have not come into contact with scabies again. If you have any questions, ask the doctor, nurse or pharmacist.

Can scabies go away without treatment?

No. And if you delay seeking treatment you risk passing the condition on to someone else.

Does having scabies affect my fertility?

No.

What happens if I get scabies when I'm pregnant or while I'm breastfeeding?

Scabies can be treated while you are pregnant or breastfeeding. However, not all types of treatment can be used. Your doctor, nurse or pharmacist will advise you. Permethrin cream is safe to use but if you are breastfeeding will need to be thoroughly washed off before any feed and re-applied if necessary.

Does scabies cause cervical cancer?

There is no evidence that scabies causes cervical cancer.

General information

How can I help protect myself from sexually transmitted infections?

It is possible to get a sexually transmitted infection by having sex with someone who has an infection. This is possible even if they have no symptoms. The following measures will help protect you from most sexually transmitted infections, including HIV, chlamydia and gonorrhoea. If you have a sexually transmitted infection they will also help prevent you from passing it on.

- Use condoms (male or female) every time you have vaginal or anal sex.
- If you have oral sex, use a condom to cover the penis, or a latex or polyurethane (soft plastic) square to cover the female genitals or male or female anus.
- If you are a woman and rub your vulva against your female partner's vulva one of you should cover the genitals with a latex or polyurethane square.
- Avoid sharing sex toys. If you do share them, wash them or cover them with a new condom before anyone else uses them.

- If you are not sure how to use condoms correctly visit www.fpa.org.uk for more information.

When should I have a test for a sexually transmitted infection?

If you, or your partner, think you may have a sexually transmitted infection it is important that you don't delay in getting a test.

You may wish to have a test if:

- you, or your partner, have or think you might have symptoms
- you have recently had unprotected sex with a new partner
- you, or your partner, have had unprotected sex with other partners
- during a vaginal examination your doctor or nurse says that the cells of the cervix are inflamed or there is a discharge
- a sexual partner tells you they have a sexually transmitted infection
- you have another sexually transmitted infection
- you are pregnant or planning a pregnancy. Don't delay seeking advice – clinics don't mind doing sexual health check-ups.

Will I know how long I've had pubic lice or scabies?

Sometimes it can be difficult to know where you

got your pubic lice or scabies from. The check-up cannot tell you how long you have had them. If you feel upset or angry about having pubic lice or scabies, don't be afraid to discuss how you feel with the staff at the clinic or general practice.

Should I tell my partner?

If the test shows that you have pubic lice or scabies then it is important that your current sexual partner(s) and any other recent partners are also checked and treated. The staff at the clinic or general practice can discuss this with you.

For additional information on scabies visit <http://www.skin-care-health.org/genital-scabies.html>